

Metrisability of Manifolds

David Gauld^{*†}

October 5, 2009

Abstract

Manifolds have uses throughout and beyond Mathematics and it is not surprising that topologists have expended a huge effort in trying to understand them. In this article we are particularly interested in the question: ‘when is a manifold metrisable?’ We describe many conditions equivalent to metrisability.

1 Introduction.

By a *manifold* is meant a connected, Hausdorff space which is locally homeomorphic to euclidean space (we take our manifolds to have no boundary). Note that because of connectedness the dimension of the euclidean space is an invariant of the manifold (unless the manifold is empty!); this is the *dimension* of the manifold. A pair (U, h) , where $U \subset M$ is open and $h : U \rightarrow \mathbb{R}^m$ is a homeomorphism, is called a *coordinate chart*.

The following notation is used. \mathbb{R} denotes the real line with the usual (order) topology while \mathbb{R}^n denotes the n th power of \mathbb{R} . \mathbb{B}^n consists of all points of \mathbb{R}^n at most 1 from the origin. The sets ω and ω_1 are, respectively, the finite and countable ordinals.

Clearly every manifold is Tychonoff. Of course manifolds share all of the local properties of euclidean space, including local compactness, local connectedness, local path or arc connectedness, first countability, local second countability, local hereditary separability, etc. As every manifold is locally compact and Hausdorff, hence completely regular, it follows that every manifold is uniformisable ([38, Proposition 11.5]). The following result shows that manifolds cannot be too big.

Proposition 1 *Let M be a (non-empty!) manifold. Then every countable subset of M is contained in an open subset which is homeomorphic to euclidean space. Hence every two points of M may be joined by an arc.*

Proof. Suppose that the dimension of M is m . Let $S \subset M$ be a countable subset, say $\langle S_n \rangle$ is such that $S = \cup_{n \geq 1} S_n$, $|S_n| = n$ and $S_n \subset S_{n+1}$.

By induction on n we choose open $V_n \subset M$ and compact $C_n \subset M$ such that

$$(i) S_n \cup C_{n-1} \subset \overset{\circ}{C}_n \text{ and } (ii) (V_n, C_n) \approx (\mathbb{R}^m, \mathbb{B}^m),$$

where $C_0 = \emptyset$.

For $n = 1$, S_1 is a singleton so V_1 may be any appropriate neighbourhood of that point while C_1 is a compact neighbourhood chosen to satisfy (ii) as well.

^{*}Supported in part by a Marsden Fund Award, UOA611, from the Royal Society of New Zealand.

[†]The author thanks Abdul Mohamad, Wang Shuquan and Frédéric Mynard for useful comments.

Suppose that V_n and C_n have been constructed. Consider

$$S = \{x \in M \mid \exists \text{ open } U \subset M \text{ with } C_n \cup \{x\} \subset U \approx \mathbb{R}^m\}.$$

S is open. S is also closed, for suppose that $z \in \bar{S} - S$. Then we may choose open $O \subset M$ with $O \approx \mathbb{R}^m$ and $z \in O$. Choose $x \in O \cap S$. Then there is open $U \subset M$ with $C_n \cup \{x\} \subset U \approx \mathbb{R}^m$. We may assume that O is small enough that $O \cap C_n = \emptyset$. Using the euclidean space structure of O we may stretch U within O so as to include z but not uncover any of C_n . Thus $z \in S$.

As M is connected and $S \neq \emptyset$ we must have $S = M$. Thus there is open $V_{n+1} \subset M$ with $S_{n+1} \cup C_n = (S_{n+1} - S_n) \cup C_n \subset V_{n+1} \approx \mathbb{R}^m$. Because $S_{n+1} \cup C_n$ is compact we may find in V_{n+1} a compact subset C_{n+1} so that (i) and (ii) hold with n replaced by $n + 1$.

Let $U_n = \overset{\circ}{C}_n$. Then U_n is open, $U_n \subset U_{n+1}$ and $U_n \approx \mathbb{R}^m$. Thus by [8], $U = \cup_{n \geq 1} U_n$ is also open with $U \approx \mathbb{R}^m$. Furthermore $S_n \subset U_n$ for each n so that $S \subset U$. ■

There has been considerable study of metrisable manifolds, especially compact manifolds. In particular it is known that there are only two metrisable manifolds of dimension 1: the circle \mathbb{S}^1 , where

$$\mathbb{S}^n = \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid x_0^2 + \dots + x_n^2 = 1\},$$

and the real line \mathbb{R} itself. In dimension 2 the compact manifolds have also been classified, this time into two sequences: the orientable manifolds, which consist of the 2-sphere \mathbb{S}^2 with n handles ($n \in \omega$) sewn on, and the non-orientable manifolds, which consist of the 2-sphere with n cross-caps ($n \in \omega - \{0\}$) sewn on. See [24, Chapter 14 and Appendix B], for example. Despite considerable progress in the study of compact manifolds in higher dimensions there has been no classification even of compact manifolds of dimension 3. Indeed, the 3-dimensional Poincaré conjecture has only now apparently been resolved after about 100 years. The original conjecture, [51], differs slightly from that posed below and was found by Poincaré to be false. Poincaré's counterexample was published in [52], where the following version was also posed. The conjecture says that if a compact manifold M of dimension 3 is such that every continuous function $\mathbb{S}^1 \rightarrow M$ extends to a continuous function $\mathbb{B}^2 \rightarrow M$, where $\mathbb{B}^2 = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}$, then M is homeomorphic to \mathbb{S}^3 . The analogue of this conjecture in dimensions higher than 3 is known to be true, [21, Corollary 7.1B] in dimension 4 and [44, Proposition B, p109] in dimension 5 and higher.

In contrast to the compact situation, where it is known that there are only countably many manifolds [14], in the nonmetrisable case there are 2^{\aleph_1} manifolds, even of dimension 2 ([47], p 669). However, there are only two nonmetrisable manifolds of dimension 1, the simpler being the open *long ray*, \mathbb{L}^+ , [9]. To ease the description of \mathbb{L}^+ we firstly give a way of constructing the positive real numbers from the non-negative integers and copies of the unit interval. Between any integer and its successor we insert a copy of the open unit interval. More precisely, let $\mathbb{R}^+ = \omega \times [0, 1) - \{(0, 0)\}$, order \mathbb{R}^+ by the lexicographic order from the natural orders on ω and $[0, 1)$ and then topologise \mathbb{R}^+ by the order topology. To get \mathbb{L}^+ we do the same thing but replace ω by ω_1 . More precisely, let $\mathbb{L}^+ = \omega_1 \times [0, 1) - \{(0, 0)\}$, order \mathbb{L}^+ by the lexicographic order from the natural orders on ω_1 and $[0, 1)$ and then topologise \mathbb{L}^+ by the order topology. The other nonmetrisable manifold of dimension 1 is the *long line*, which is obtained by joining together two copies of \mathbb{L}^+ at their $(0, 0)$ ends in much the same way as one may reconstruct the real line \mathbb{R} by joining together two copies of $(0, \infty)$ at their 0 ends, thinking of one copy as giving the positive reals and the other the negative reals. More precisely, let \mathbb{L} be the disjoint union of two copies of \mathbb{L}^+ (call them \mathbb{L}^+ and \mathbb{L}^- respectively, with ordering $<^+$ and $<^-$ respectively) as well as a single point, which we denote by 0, order \mathbb{L} by $x < y$ when $x <^+ y$ in \mathbb{L}^+ , when $x = 0$ and $y \in \mathbb{L}^+$, when $x \in \mathbb{L}^-$ and $y \in \mathbb{L}^+$ or $y = 0$ and when $x, y \in \mathbb{L}^-$ and $y <^- x$, and topologise \mathbb{L} by the order topology.

The survey articles [47] and [49] are good sources of information about nonmetrisable manifolds.

A significant question in topology is that of deciding when a topological space is metrisable, there being many criteria which have now been developed to answer the question. Perhaps the most natural is the following: a topological space is metrisable if and only if it is paracompact, Hausdorff and locally metrisable, see [58] and [37, Theorem 2.68]. Note that manifolds are always Hausdorff and locally metrisable so this criterion gives a criterion for the metrisability of a manifold, viz that a manifold is metrisable if and only if it is paracompact. Many metrisation criteria have been discovered for manifolds, as seen by Theorem 2 below, which lists criteria which require at least some of the extra properties possessed by manifolds. Of course one must not be surprised if conditions which in general topological spaces are considerably weaker than metrisability are actually equivalent to metrisability in the presence of the extra topological conditions which always hold for a manifold: such a condition is that of being nearly meta-Lindelöf, 10 in Theorem 2 below. Similarly one should not be surprised to find conditions which are normally stronger than metrisability: such a condition is that M may be embedded in euclidean space, 35 in Theorem 2. Finally one may expect to find conditions which in a general topological space have no immediate connection with metrisability: such a condition is second countability, 26 in Theorem 2.

2 Definitions.

In this section we list numerous definitions relevant to the question of metrisability..

Definitions: Let X be a topological space and \mathcal{F} a family of subsets of X . Then:

- X is *submetrisable* if there is a metric topology on X which is contained in the given topology;
- X is *Polish* if X is a separable, complete metric space;
- X is *paracompact* (respectively *metacompact*, *paraLindelöf* and *metaLindelöf*) if every open cover \mathcal{U} has a locally finite (respectively point finite, locally countable, and point countable) open refinement, ie there is another open cover \mathcal{V} such that each member of \mathcal{V} is a subset of some member of \mathcal{U} and each point of X has a neighbourhood meeting only finitely (respectively lies in only finitely, has a neighbourhood meeting only countably, and lies in only countably) many members of \mathcal{V} ;
- X is *finitistic* (respectively *strongly finitistic*) if every open cover of X has an open refinement \mathcal{V} and there is an integer m such that each point of X lies in (respectively has a neighbourhood which meets) at most m members of \mathcal{V} (finitistic spaces have also been called *boundedly metacompact* and strongly finitistic spaces have also been called *boundedly paracompact*);
- X is *strongly paracompact* if every open cover \mathcal{U} has a star-finite open refinement \mathcal{V} , ie for any $V \in \mathcal{V}$ the set $\{W \in \mathcal{V} / V \cap W \neq \emptyset\}$ is finite. If in addition, given \mathcal{U} , there is an integer m such that $\{W \in \mathcal{V} / V \cap W \neq \emptyset\}$ contains at most m members then X is *star finitistic*;
- X is *screenable* (respectively *σ -metacompact* and *σ -paraLindelöf*) if every open cover \mathcal{U} has an open refinement \mathcal{V} which can be decomposed as $\mathcal{V} = \cup_{n \in \omega} \mathcal{V}_n$ such that each \mathcal{V}_n is disjoint (respectively point finite and locally countable);

- X is (linearly) $[\omega_1]$ -Lindelöf if every open cover (which is a chain) [which has cardinality ω_1] has a countable subcover;
- X is (nearly) [linearly ω_1]-metaLindelöf if every open cover \mathcal{U} of X [with $|\mathcal{U}| = \omega_1$] has an open refinement which is point-countable (on a dense subset);
- X is almost metaLindelöf if for every open cover \mathcal{U} there is a collection \mathcal{V} of open subsets of X such that each member of \mathcal{V} lies in some member of \mathcal{U} , that each point of X lies in at most countably many members of \mathcal{V} , and that $X = \bigcup \{\overline{V} / V \in \mathcal{V}\}$.
- X is (strongly) hereditarily Lindelöf if every subspace (of the countably infinite power) of X is Lindelöf;
- X is k -Lindelöf provided every open k -cover (ie every compact subset of X lies in some member of the cover) has a countable k -subcover;
- X is (strongly) hereditarily separable if every subspace (of the countably infinite power) of X is separable;
- X is Hurewicz if for each sequence $\langle \mathcal{U}_n \rangle$ of open covers of X there is a sequence $\langle \mathcal{V}_n \rangle$ such that \mathcal{V}_n is a finite subset of \mathcal{U}_n for each $n \in \omega$ and $\bigcup_{n \in \omega} \mathcal{V}_n$ covers X (note the alternative definition of Hurewicz, [17]: X is Hurewicz if for each sequence $\langle \mathcal{U}_n \rangle$ of open covers of X there is a sequence $\langle \mathcal{V}_n \rangle$ such that \mathcal{V}_n is a finite subset of \mathcal{U}_n and for each $x \in X$ we have $x \in \bigcup \mathcal{V}_n$ for all but finitely many $n \in \omega$. For a manifold these two conditions are equivalent.);
- X is selectively screenable, [1], if for each sequence $\langle \mathcal{U}_n \rangle$ of open covers of X there is a sequence $\langle \mathcal{V}_n \rangle$ such that \mathcal{V}_n is a family of pairwise disjoint open sets refining \mathcal{U}_n for each $n \in \omega$ and $\bigcup_{n \in \omega} \mathcal{V}_n$ covers X ;
- X is hemicompact if there is an increasing sequence $\langle K_n \rangle$ of compact subsets of X such that for any compact $K \subset X$ there is n such that $K \subset K_n$;
- X is cosmic if there is a countable family \mathcal{C} of closed subsets of X such that for each point $x \in X$ and each open set U containing x there is a set $C \in \mathcal{C}$ such that $x \in C \subset U$;
- X is an \aleph_0 -space ([33, page 493]) provided that it has a countable k -network, i.e. a countable collection \mathcal{N} such that if $K \subset U$ with K compact and U open then $K \subset N \subset U$ for some $N \in \mathcal{N}$;
- X is an \aleph -space ([33, page 493]) provided that it has a σ -locally finite k -network;
- X has the Moving Off Property, [35], provided that every family \mathcal{K} of non-empty compact subsets of X large enough to contain for each compact $C \subset X$ a disjoint $K \in \mathcal{K}$ has an infinite subfamily with a discrete open expansion;
- X is a q -space if each point admits a sequence of neighbourhoods Q_n such that $x_n \in Q_n$ implies that $\langle x_n \rangle$ clusters;
- X is Fréchet or Fréchet-Urysohn if whenever $x \in \overline{A}$ there is a sequence $\langle x_n \rangle$ in A that converges to x ;
- X is a k -space if A is closed whenever $A \cap K$ is closed for every compact subset $K \subset X$;
- X is Lašnev if it is the image of a metrisable space under a closed map;

- X is *analytic* if it is the continuous image of a Polish space (equivalently of the irrational numbers);
- X is M_1 if it has a σ -closure preserving base (ie a base \mathcal{B} such that there is a decomposition $\mathcal{B} = \cup_{n=1}^{\infty} \mathcal{B}_n$ where for each n and each $\mathcal{F} \subset \mathcal{B}_n$ we have $\overline{\cup \mathcal{F}} = \cup \{\bar{F} / F \in \mathcal{F}\}$);
- X is *stratifiable* or M_3 if there is a function G which assigns to each $n \in \omega$ and closed set $A \subset X$ an open set $G(n, A)$ containing A such that $A = \cap_n \overline{G(n, A)}$ and if $A \subset B$ then $G(n, A) \subset G(n, B)$;
- X is *perfectly normal* if for every pair A, B of disjoint closed subsets of X there is a continuous function $f : X \rightarrow \mathbb{R}$ such that $f^{-1}(0) = A$ and $f^{-1}(1) = B$;
- X is *monotonically normal* if for each open $U \subset X$ and each $x \in U$ it is possible to choose an open set $\mu(x, U)$ such that $x \in \mu(x, U) \subset U$ and such that if $\mu(x, U) \cap \mu(y, V) \neq \emptyset$ then either $x \in V$ or $y \in U$;
- X is *extremely normal* if for each open $U \subset X$ and each $x \in U$ it is possible to choose an open set $\nu(x, U)$ such that $x \in \nu(x, U) \subset U$ and such that if $\nu(x, U) \cap \nu(y, V) \neq \emptyset$ and $x \neq y$ then either $\nu(x, U) \subset V$ or $\nu(y, V) \subset U$;
- X is *weakly normal* if for every pair A, B of disjoint closed subsets of X there is a continuous function $f : X \rightarrow S$, for some separable metric space S , such that $f(A) \cap f(B) = \emptyset$;
- X is a *Moore space* if it is regular and has a development, ie a sequence $\langle \mathcal{U}_n \rangle$ of open covers such that for each $x \in X$ the collection $\{st(x, \mathcal{U}_n) : n \in \omega\}$ forms a neighbourhood basis at x ;
- X has a *regular G_δ -diagonal* if the diagonal Δ is a *regular G_δ -subset* of X^2 , ie there is a sequence $\langle U_n \rangle$ of open subsets of X^2 such that $\Delta = \cap U_n = \overline{\cap U_n}$.
- X has a *quasi-regular G_δ -diagonal* if there is a sequence $\langle U_n \rangle$ of open subsets of X^2 such that for each $(x, y) \in X^2 - \Delta$ there is n with $(x, x) \in U_n$ but $(x, y) \notin \overline{U_n}$.
- X has a *G_δ^* -diagonal* if there is a sequence $\langle \mathcal{G}_n \rangle$ of open covers of X such that for each $x, y \in X$ with $x \neq y$ there is n with $st(x, \mathcal{G}_n) \subset X - \{y\}$.
- X has a *quasi- G_δ^* -diagonal* if there is a sequence $\langle \mathcal{G}_n \rangle$ of families of open subsets of X such that for each $x, y \in X$ with $x \neq y$ there is n with $x \in st(x, \overline{\mathcal{G}_n}) \subset X - \{y\}$.
- X is *θ -refinable* if every open cover can be refined to an open θ -cover, ie a cover \mathcal{U} which can be expressed as $\cup_{n \in \omega} \mathcal{U}_n$ where each \mathcal{U}_n covers X and for each $x \in X$ there is n such that $ord(x, \mathcal{U}_n) < \omega$;
- X is *subparacompact* if every open cover has a σ -discrete closed refinement;
- X has *property pp*, [42], provided that each open cover \mathcal{U} of X has an open refinement \mathcal{V} such that for each choice function $f : \mathcal{V} \rightarrow X$ with $f(V) \in V$ for each $V \in \mathcal{V}$ the set $f(\mathcal{V})$ is closed and discrete in X ;
- X has *property (a)*, [42], provided that for each open cover \mathcal{U} of X and each dense subset $D \subset X$ there is a subset $C \subset D$ such that C is closed and discrete in X and $st(C, \mathcal{U}) = X$;

- X has a *base of countable order*, \mathcal{B} , if whenever $\mathcal{C} \subset \mathcal{B}$ is a collection such that each member of \mathcal{C} contains a particular point $p \in X$ and for each $C \in \mathcal{C}$ there is $D \in \mathcal{C}$ with D a proper subset of C then \mathcal{C} is a local base at p ;
- X is *pseudocomplete* provided that it has a sequence $\langle \mathcal{B}_n \rangle$ of π -bases ($\mathcal{B} \subset 2^X - \{\emptyset\}$ is a π -base if every non-empty open subset of X contains some member of \mathcal{B}) such that if $B_n \in \mathcal{B}_n$ and $\overline{B_{n+1}} \subset B_n$ for each n , then $\bigcap_{n \in \omega} B_n \neq \emptyset$;
- X has the *countable chain condition* (abbreviated *ccc*) if every pairwise disjoint family of open subsets is countable;
- X is *countably tight* if for each $A \subset X$ and each $x \in \bar{A}$ there is a countable $B \subset A$ for which $x \in \bar{B}$;
- X is *countably fan tight* if whenever $x \in \bigcap_{n \in \omega} \overline{A_n}$ there are finite sets $B_n \subset A_n$ such that $x \in \overline{\bigcup_{n \in \omega} B_n}$;
- X is *countably strongly fan tight* if whenever $x \in \bigcap_{n \in \omega} \overline{A_n}$ there is a sequence $\langle a_n \rangle$ such that $a_n \in A_n$ for each n and $x \in \overline{\{a_n / n \in \omega\}}$;
- X is *sequential* if for each $A \subset X$, the set A is closed whenever for each sequence of points of A each limit point is also in A ;
- X is *weakly α -favourable* if there is a winning strategy for player α in the Banach-Mazur game (defined below);
- X is *strongly α -favourable* if there is a stationary winning strategy for player α in the Choquet game (defined below);
- for each $x \in X$ the *star* of x in \mathcal{F} is $st(x, \mathcal{F}) = \bigcup \{F \in \mathcal{F} : x \in F\}$;
- X is *Baire* provided that the intersection of any countable collection of dense G_δ subsets is dense;
- X is *Volterra*, [29], provided that the intersection of any two dense G_δ subsets is dense;
- X is *strongly Baire* provided that X is regular and there is a dense subset $D \subset X$ such that β does not have a winning strategy in the game $\mathcal{G}_S(D)$ played on X .
- \mathcal{F} is *point-star-open* if for each $x \in X$ the set $st(x, \mathcal{F})$ is open.
- The *Banach-Mazur game* has two players α and β whose play alternates. Player β begins by choosing a non-empty open subset of X . After that the players choose successive non-empty open subsets of their opponent's previous move. *Player α wins* iff the intersection of the sets is non-empty; otherwise *player β wins*.
- The *Choquet game* has two players α and β whose play alternates. Player β begins by choosing a point in an open subset of X , say $x_0 \in V_0 \subset X$. After that the players alternate with α choosing an open set $U_n \subset X$ with $x_n \in U_n \subset V_n$ then β chooses a point x_{n+1} and an open set V_{n+1} with $x_{n+1} \in V_{n+1} \subset U_n$. *Player α wins* iff the intersection of the sets is non-empty; otherwise *player β wins*.

- Gruenhage's game $G_{K,L}^o(X)$, [34], has, at the n^{th} stage, player K choose a compactum $K_n \subset X$ after which player L chooses another compactum $L_n \subset X$ so that $L_n \cap K_i = \emptyset$ for each $i \leq n$. *Player K wins* if $\langle L_n \rangle_{n \in \omega}$ has a discrete open expansion, ie there is a sequence $\langle U_n \rangle_{n \in \omega}$ of open sets such that $L_n \subset U_n$ and $\forall x \in X, \exists U \subset M$ open such that $x \in U$ and U meets at most one of the sets U_n .
- For a dense subset $D \subset X$ the game $\mathcal{G}_S(D)$ has two players α and β whose play alternates. Player β begins by choosing a non-empty open subset V_n of X . After that the players choose successive non-empty open subsets of their opponent's previous move, β choosing sets V_n and α choosing sets U_n . *Player α wins* iff the intersection of the sets is non-empty and each sequence $\langle x_n \rangle$, for which $x_n \in U_n \cap D$, clusters in X ; otherwise *player β wins*.
- For an ordinal k and families \mathbb{A} and \mathbb{B} of collections of subsets of a space X let $\mathcal{G}_c^k(\mathbb{A}, \mathbb{B})$ be the game played as follows, [6]: at the l^{th} stage of the game, $l < k$, Player One chooses a member $\mathcal{A}_l \in \mathbb{A}$ then Player Two chooses a pairwise disjoint family \mathcal{T}_l which refines \mathcal{A}_l . The play $\mathcal{A}_0, \mathcal{T}_0, \dots, \mathcal{A}_l, \mathcal{T}_l, \dots$ $l < k$ is won by Player Two provided that $\cup_{l < k} \mathcal{T}_l \in \mathbb{B}$; otherwise Player One wins. The game $\mathcal{G}_c^\omega(\mathbb{A}, \mathbb{B})$ denoted by $\mathcal{G}_c(\mathbb{A}, \mathbb{B})$.
- When players α and β play a topological game, a *strategy* for α is a function which tells α what points or sets to select given all the previous points and sets chosen by β . A *stationary strategy* for α is a function which tells α what points or sets to select given only the most recent choice of points and sets chosen by β . A *winning (stationary) strategy* for α is a (stationary) strategy which guarantees that α will win whatever moves β might make.

We will denote by $C_k(X, Y)$ (respectively $C_p(X, Y)$) the space of all continuous functions from X to Y with the compact-open topology (respectively the topology of pointwise convergence).

We will denote by $\$$ the space $\{0, 1\}$ with the Sierpinski topology $\{\emptyset, \$, \{0\}\}$. Then for any space X we denote by $[X, \$]$ the space of continuous functions from X to $\$$ with the *upper Kuratowski topology*, i.e. that in which a subset $\mathcal{F} \subset [X, \$]$ is open if and only if

- (i) for each $f \in \mathcal{F}$ and each $g \in [X, \$]$ if $g \leq f$ then $g \in \mathcal{F}$;
- (ii) if $\mathcal{G} \subset [X, \$]$ is such that $\inf \mathcal{G} \in \mathcal{F}$ then there is a finite subfamily $\mathcal{G}' \subset \mathcal{G}$ with $\inf \mathcal{G}' \in \mathcal{F}$.

In this definition we are using the usual ordering on $\{0, 1\}$ when discussing \leq and \inf . Of course identifying a closed subset of X with its characteristic function gives a bijective correspondence between $[X, \$]$ and the collection of closed subsets of X . This topology is also variously known as the *cocompact topology* and the *upper Fell topology*, especially when looked at as a topology on the set 2^X of non-empty closed subsets of X . Letting $U^+ = \{C \in 2^X / C \subset U\}$ for $U \subset X$, this topology has as subbasis $\{U^+ / U \text{ is open in } X \text{ and } X - U \text{ is compact}\}$. The *Fell topology*, [19], denoted by τ_{fell} , has as subbasis

$$\{U^+ / U \text{ is open in } X \text{ and } X - U \text{ is compact}\} \cup \{U^- / U \text{ is open in } X\},$$

where $U^- = \{C \in 2^X / C \cap U \neq \emptyset\}$.

3 Criteria for metrisability.

We now state and outline the proof of the main theorem. It is believed that no two conditions are equivalent in a general topological space, though, as will be noticed at the start of the proof of Theorem 2, there may be a chain of implications holding in a general space.

Theorem 2 *Let M be a manifold. Then the following are equivalent:*

1. M is metrisable;
2. M is paracompact;
3. M is strongly paracompact;
4. M is screenable;
5. M is metacompact;
6. M is σ -metacompact;
7. M is paraLindelöf;
8. M is σ -paraLindelöf;
9. M is metaLindelöf;
10. M is nearly metaLindelöf;
11. M is Lindelöf;
12. M is linearly Lindelöf;
13. M is ω_1 -Lindelöf;
14. M is ω_1 -metaLindelöf;
15. M is nearly linearly ω_1 -metaLindelöf;
16. M is almost metaLindelöf;
17. M is hereditarily Lindelöf;
18. M is strongly hereditarily Lindelöf;
19. M is k -Lindelöf;
20. M is an \aleph_0 -space;
21. M is cosmic;
22. M is an \aleph -space;
23. M has a star-countable k -network;
24. M has a point-countable k -network;
25. M has a k -network which is point-countable on some dense subset of M ;
26. M is second countable;
27. M is hemicompact;
28. M is σ -compact;
29. M is Hurewicz;

30. M satisfies the selection criterion $S_1(\mathcal{K}, \Gamma)$: for each sequence $\langle \mathcal{U}_n \rangle$ of open k -covers of X there is a sequence $\langle U_n \rangle$ with $U_n \in \mathcal{U}_n$ for each n , infinitely many of the sets U_n are distinct and each finite subset of X lies in all but finitely many of the sets U_n ;
31. Player Two has a winning strategy in the game $G_c^{n+1}(\emptyset, \emptyset)$ played on M , where \emptyset denotes the family of all open covers of M ;
32. Player Two has a winning strategy in the game $G_c(\emptyset, \emptyset)$ played on M , where \emptyset denotes the family of all open covers of M ;
33. M is selectively screenable;
34. only countably many coordinate charts are needed to cover M ;
35. M may be embedded in some euclidean space;
36. M may be embedded properly in some euclidean space;
37. M is completely metrisable;
38. there is a continuous discrete map $f : M \rightarrow X$ where X is Hausdorff and second countable;
39. M is Lašnev;
40. M is an M_1 -space;
41. M is stratifiable;
42. M is finitistic;
43. M is strongly finitistic;
44. M is star finitistic;
45. there is an open cover \mathcal{U} of M such that for each $x \in M$ the set $st(x, \mathcal{U})$ is homeomorphic to an open subset of \mathbb{R}^m ;
46. there is a point-star-open cover \mathcal{U} of M such that for each $x \in M$ the set $st(x, \mathcal{U})$ is Lindelöf;
47. there is a point-star-open cover \mathcal{U} of M such that for each $x \in M$ the set $st(x, \mathcal{U})$ is metrisable;
48. the tangent microbundle on M is equivalent to a fibre bundle;
49. M is a normal Moore space;
50. M is a normal θ -refinable space;
51. M is a normal subparacompact space;
52. M is a normal space which has a σ -discrete cover by compact subsets;
53. $M \times M$ is perfectly normal;
54. M is a normal space which has a sequence $\langle \mathcal{U}_n \rangle_{n \in \omega}$ of open covers with $\bigcap_n \overline{st(x, \mathcal{U}_n)} = \{x\}$ for each $x \in M$;


55. M is perfectly normal and there is a sequence $\langle \mathcal{U}_n \rangle_{n \in \omega}$ of families of open sets such that $\bigcap_{n \in C(x)} \overline{st(x, \mathcal{U}_n)} = \{x\}$ for each $x \in M$, where
- $$C(x) = \{n \in \omega \mid \exists U \in \mathcal{U}_n \text{ with } x \in U\};$$
56. M is separable and there is a sequence $\langle \mathcal{C}_n \rangle_{n \in \omega}$ of point-star-open covers such that $\bigcap_n \overline{st(x, \mathcal{C}_n)} = \{x\}$ for each $x \in M$ and for each $x, y \in M$ and each $n \in \omega$ we have $y \in \overline{st(x, \mathcal{C}_n)}$ if and only if $x \in \overline{st(y, \mathcal{C}_n)}$;
57. M is separable and there is a sequence $\langle \mathcal{C}_n \rangle_{n \in \omega}$ of point-star-open covers such that $\bigcap_n \overline{st(x, \mathcal{C}_n)} = \{x\}$ for each $x \in M$ and for each $x \in M$ and each $n \in \omega$, $\text{ord}(x, \mathcal{C}_n)$ is finite;
58. M is separable and hereditarily normal and there is a sequence $\langle \mathcal{C}_n \rangle_{n \in \omega}$ of point-star-open covers such that $\bigcap_n \overline{st(x, \mathcal{C}_n)} = \{x\}$ for each $x \in M$;
59. M is separable and there is a sequence $\langle \mathcal{U}_n \rangle_{n \in \omega}$ of families of open sets such that $\bigcap_{n \in C(x)} \overline{st(x, \mathcal{U}_n)} = \{x\}$ for each $x \in M$, and $\text{ord}(x, \mathcal{C}_n)$ is countable for each $x \in M$ and each $n \in \omega$;
60. $M \times M$ has a countable sequence $\langle U_n : n \in \omega \rangle$ of open subsets, such that for all $(x, y) \in M \times M - \Delta$, there is $n \in \omega$ such that $(x, x) \in U_n$ but $(x, y) \notin \overline{U_n}$;
61. For every subset $A \subset M$ there is a continuous injection $f : M \rightarrow Y$, where Y is a metrisable space, such that $f(A) \cap f(M - A) = \emptyset$;
62. For every subset $A \subset M$ there is a continuous $f : M \rightarrow Y$, where Y is a space with a quasi-regular- G_δ -diagonal, such that $f(A) \cap f(M - A) = \emptyset$;
63. M is weakly normal with a G_δ^* -diagonal;
64. M has a quasi- G_δ^* -diagonal and for every closed subset $A \subset M$ there is a countable family \mathcal{G} of open subsets such that, for every $x \in A$ and $y \in X - A$, there is a $G \in \mathcal{G}$ with $x \in G, y \notin \overline{G}$;
65. M has a regular G_δ -diagonal;
66. M is submetrisable;
67. M is separable and monotonically normal;
68. $M \times M$ is monotonically normal;
69. M is monotonically normal and of dimension ≥ 2 or $M \approx \mathbb{S}^1$ or \mathbb{R} ;
70. M is extremely normal;
71. M has the Moving Off Property;
72. M has property pp;
73. every open cover of M has an open refinement \mathcal{V} such that for every choice function $f : \mathcal{V} \rightarrow M$ the set $f(\mathcal{V})$ is closed in M ;
74. every open cover of M has an open refinement \mathcal{V} such that for every choice function $f : \mathcal{V} \rightarrow M$ the set $f(\mathcal{V})$ is discrete in M ;

75. M is a point-countable union of open subspaces each of which is metrisable;
76. M has a point-countable basis;
77. M is separable and M^ω is a countable union of metrisable subspaces;
78. $C_k(M, \mathbb{R})$ is Polish;
79. $C_k(M, \mathbb{R})$ is completely metrisable;
80. $C_k(M, \mathbb{R})$ is first countable;
81. $C_k(M, \mathbb{R})$ is second countable;
82. $C_k(M, \mathbb{R})$ is a q -space;
83. $C_k(M, \mathbb{R})$ is Fréchet;
84. $C_k(M, \mathbb{R})$ is countably tight;
85. $C_k(M)$ has countable strong fan tightness;
86. $C_k(M, \mathbb{R})$ is an \aleph_0 -space;
87. $C_k(M, \mathbb{R})$ is cosmic;
88. $C_k(M, \mathbb{R})$ is analytic;
89. $C(M, \mathbb{R})$ satisfies the selection criterion $S_1(\Omega_{\underline{0}}^k, \Sigma_{\underline{0}}^p)$: for each sequence $\langle F_n \rangle$ of subsets of $C(M, \mathbb{R})$ whose compact-open closures contain the constant function $\underline{0}$ there is a sequence $\langle f_n \rangle$, infinitely many members of which are distinct, with $f_n \in F_n$ for all n and $\langle f_n \rangle$ converges pointwise to $\underline{0}$;
90. $C_p(M, \mathbb{R})$ has countable tightness;
91. $C_p(M, \mathbb{R})$ has countable fan tightness;
92. $C_p(M, \mathbb{R})$ is analytic;
93. $C_p(M, \mathbb{R})$ is hereditarily separable;
94. $C_p(M, \mathbb{R})$ (equivalently $C_k(M, \mathbb{R})$) is separable;
95. $[M, \$]$ is first countable;
96. $[M, \$]$ is countably tight;
97. $[M, \$]$ is sequential;
98. $(2^X, \tau_{\text{fell}})$ is metrisable;
99. $(2^X, \tau_{\text{fell}})$ is countably tight;
100. $(2^X, \tau_{\text{fell}})$ is sequential;
101. $(2^X, \tau_{\text{fell}})$ is radial;
102. K has a winning strategy in Gruenhage's game $G_{K,L}^o(M)$;

103. $C_k(M, \mathbb{R})$ is strongly α -favourable;
104. $C_k(M, \mathbb{R})$ is weakly α -favourable;
105. $C_k(M, \mathbb{R})$ is pseudocomplete;
106. $C_k(M, \mathbb{R})$ is strongly Baire;
107. $C_k(M, \mathbb{R})$ is Baire;
108. $C_k(M, \mathbb{R})$ is Volterra.

Outline of the proof of theorem 2.

The following diagram shows how items 1-34 are related:



$\boxed{\mathbf{m}}$ =metrisable; $\boxed{\mathbf{sc}}$ =second countable; $\boxed{\mathbf{shL}}$ =strongly hereditarily Lindelöf; $\boxed{\mathbf{hL}}$ =hereditarily Lindelöf; $\boxed{\sigma\mathbf{c}}$ = σ -compact; $\boxed{\mathbf{hc}}$ =hemicompact; $\boxed{\mathbf{H}}$ =Hurewicz; $\boxed{\mathbf{SKT}}$ =satisfies $S_1(\mathcal{K}, \Gamma)$; $\boxed{\mathbf{cch}}$ =countably many charts cover; $\boxed{\mathbf{N}_0}$ = \mathbf{N}_0 -space; $\boxed{\mathbf{N}}$ = \mathbf{N} -space; $\boxed{\mathbf{skn}}$ =has a star-countable k -network; $\boxed{\mathbf{pkn}}$ =has a point-countable k -network; $\boxed{\mathbf{npkn}}$ =has a k -network which is point-countable on a dense subset; $\boxed{\mathbf{kL}}$ = k -Lindelöf; $\boxed{\mathbf{c}}$ =cosmic; $\boxed{\mathbf{L}}$ =Lindelöf; $\boxed{\mathbf{IL}}$ =linearly Lindelöf; $\boxed{\omega_1\mathbf{L}}$ = ω_1 -Lindelöf; $\boxed{\mathbf{spc}}$ =strongly paracompact; $\boxed{\mathbf{pc}}$ =paracompact; $\boxed{\mathbf{mc}}$ =metacompact; $\boxed{\mathbf{s}}$ =screenable; $\boxed{\mathbf{ss}}$ =selectively screenable; $\boxed{\mathbf{pL}}$ =paraLindelöf; $\boxed{\sigma\mathbf{mc}}$ = σ -metacompact; $\boxed{\sigma\mathbf{pL}}$ = σ -paraLindelöf; $\boxed{\mathbf{mL}}$ =metaLindelöf; $\boxed{\mathbf{amL}}$ =almost metaLindelöf; $\boxed{\mathbf{nmL}}$ =nearly

metaLindelöf; $\boxed{\omega_1\text{mL}} = \omega_1\text{-metaLindelöf}$; $\boxed{\text{nl}\omega_1\text{mL}} = \text{nearly linearly } \omega_1\text{-metaLindelöf}$; $\boxed{G(n+1)}$ = Player Two has a winning strategy in the game $G_c^{n+1}(\emptyset, \emptyset)$; $\boxed{G\omega}$ = Player Two has a winning strategy in the game $G_c(\emptyset, \emptyset)$.

All arrows denote implications. Downward sloping arrows show an implication which holds in an arbitrary topological space. Upward sloping arrows require one or more properties of manifolds to realise the implication. $\boxed{\text{mL}} \Rightarrow \boxed{L}$ in every locally separable and connected space. $\boxed{\text{amL}} \Rightarrow \boxed{L}$ in every regular, locally separable and connected space, [27]. $\boxed{\text{nmL}} \Rightarrow \boxed{\text{mL}}$ in every locally hereditarily separable space. $\boxed{L} \Rightarrow \boxed{\text{spc}}$ in every T_3 space. $\boxed{\omega_1 L} \Rightarrow \boxed{L}$ in every locally metrisable space, [5]. $\boxed{L} \Rightarrow \boxed{\text{sc}}$ in every locally second countable space. $\boxed{L} \Rightarrow \boxed{\text{hc}}$ in every locally compact space. $\boxed{\text{cch}} \Rightarrow \boxed{L}$ because a countable union of Lindelöf sets is Lindelöf. $\boxed{\text{sc}} \Rightarrow \boxed{\text{m}}$ in every T_3 space (Urysohn's metrisation theorem). $\boxed{\text{m}} \Rightarrow \boxed{G(n+1)}$ in every space having covering dimension at most n , [6, Thm 2.4]. $\boxed{\omega_1\text{mL}} \Rightarrow \boxed{\text{mL}}$ in every locally second countable space, [31]. $\boxed{\text{nl}\omega_1\text{mL}} \Rightarrow \boxed{\omega_1\text{mL}}$ in every locally hereditarily separable space, [31]. $\boxed{\text{pkn}} \Rightarrow \boxed{\text{mL}}$ in every regular Fréchet space. $\boxed{\text{npkn}} \Rightarrow \boxed{\text{pkn}}$ in every regular, locally compact, locally hereditarily separable space.

By [50, Proposition 7.3.9] we conclude that a metrisable n -manifold, being separable and of covering dimension n , embeds in \mathbb{R}^{2n+1} , so $1 \Rightarrow 35$. By choosing a proper continuous real-valued function on M we can add a further coordinate to embed M in \mathbb{R}^{2n+2} so that the image is closed, ie the embedding is proper, hence $1 \Rightarrow 36$. It is clear that $36 \Rightarrow 37$.

Every second countable Hausdorff space satisfies 38 so $26 \Rightarrow 38$. Conversely, given the situation of 38, if \mathcal{B} is a countable base for the topology on X then the Poincaré-Volterra Lemma of [20, Lemma 23.2] asserts that

$$\{U \subset M \mid U \text{ is second countable and} \\ \text{there is } V \in \mathcal{B} \text{ such that } U \text{ is a component of } f^{-1}(V)\}$$

is a countable base for M .

Clearly every metrisable space is Lašnev so $1 \Rightarrow 39$. The implication $39 \Rightarrow 40$ is [33, Theorem 5.5]. It is easy to show that $40 \Rightarrow 41$. The implication $41 \Rightarrow 2$ is [33, Theorem 5.7].

The conditions 1, 42, 43 and 44 are shown to be equivalent in [16].

The equivalence of conditions 1 and 45-48 is established as follows: $1 \Rightarrow 45$ is reasonably straightforward making use of the fact that metrisable manifolds are σ -compact. Then $45 \Rightarrow 46$ is trivial. $46 \Rightarrow 47$ requires use of Urysohn's metrisation theorem to deduce that the Lindelöf stars are metrisable. $47 \Rightarrow 11$ requires some delicate manoeuvres; see [28]. $48 \Rightarrow 45$ is also found in [28] while $1 \Rightarrow 48$ is [40, Corollary 2].

The implication $1 \Rightarrow 49$ holds in every topological space while its converse holds provided that the space is locally compact and locally connected, [53] or [54, Theorem 3.4]. The equivalence of 49 and 50 comes from [60, Theorem 3], while the equivalence of 49, 50, 51 and 52 is discussed in [48, Theorem 8.2].

The equivalence of conditions 1, 53 and 54 is referred to briefly in [25]. The implications $1 \Rightarrow 53 \Rightarrow 54$ hold in any topological space and the implication $54 \Rightarrow 1$ uses some properties of a manifold.

The equivalence of conditions 1 and 55-58 is discussed in [45].

Proofs of the equivalence of 1 and 59 may be found in [23] and of 1 and 60-64 may be found in [22].

The implication $65 \Rightarrow 1$ holds in every locally compact, locally connected space ([33, Theorem 2.15(b)]) and, as noted in [33, p. 430], every submetrisable space has a regular G_δ -diagonal so $66 \Rightarrow 65$.

Every metric space is monotonically normal and every metrisable manifold is second countable, hence separable, so $1 \Rightarrow 67$. To get the converse implication $67 \Rightarrow 2$ use is made of the fact

that every monotonically normal space is hereditarily collectionwise normal ([36]), and hence no separable monotonically normal space contains a copy of ω_1 . On the other hand in [7, Theorem I] it is shown that a monotonically normal space is paracompact if and only if it does not contain a stationary subset of a regular uncountable ordinal.

If M is metrisable, so is $M \times M$, so that $M \times M$ is monotonically normal and hence $1 \Rightarrow 68$. The converse follows from a metrisability result of [36] as manifolds are locally countably compact.

The criterion 69 is [7, corollary 2.3(e)], except that we have listed all of the metrisable 1-manifolds.

Every metrisable space is extremely normal. The implication $70 \Rightarrow 2$ is found in [61].

The equivalence of conditions 1, 71 and 107 is discussed in [10].

It is readily shown that every T_1 -space which is paracompact has property pp. We now obtain the implication $72 \Rightarrow 5$. Suppose that \mathcal{U} is an open cover of M . Use the property pp to find an open refinement \mathcal{V} such that for each choice function $f : \mathcal{V} \rightarrow M$ with $f(V) \in V$ for each $V \in \mathcal{V}$ the set $f(\mathcal{V})$ is closed and discrete. We will show that \mathcal{V} is point-finite. Suppose to the contrary that $x \in M$ is such that $\{V \in \mathcal{V} / x \in V\}$ is infinite; let $\langle V_n \rangle$ be a sequence of distinct members of \mathcal{V} each of which contains x . Because M is a manifold, hence first countable, we may choose a countable neighbourhood basis $\{W_n / n \in \omega\}$ at x . Note that for each n , $V_n \cap W_n - \{x\} \neq \emptyset$ as M has no isolated points. Choose a function $f : \mathcal{V} \rightarrow M$ as follows: if $V \in \mathcal{V}$ but $V \neq V_n$ for each n then choose $f(V) \in V - \{x\}$ arbitrarily; if $V = V_n$ choose $f(V_n) \in V_n \cap W_n - \{x\}$. Then $x \in \overline{f(\mathcal{V})} - f(\mathcal{V})$ so that $f(\mathcal{V})$ is not closed, contrary to the choice of \mathcal{V} . Thus \mathcal{V} is point-finite so M is metacompact.

It is easy to show that conditions 73 and 74 are equivalent to each other, and hence also to 72; cf [26, Lemma 2.3].

Details for the implication $75 \Rightarrow 9$ appear in [28], while details for the implication $76 \Rightarrow 1$ appear in [18]. Of course $26 \Rightarrow 76$.

The implication $77 \Rightarrow 1$ is a consequence of the more general result that if the countable power of a topological space X is a countable union of metrisable subspaces and in X discrete families of open sets are countable then X is metrisable, [59].

The equivalence of conditions 1 and 78 to 97, excluding 80, 85 and 89, is shown in [30]. A number of properties of manifolds are required, including that every manifold is a q -space and a k -space, and some of the equivalences to metrisability already proved.

Conditions 80 and 85 are shown to be equivalent to condition 11 in [13, Theorem 6] using Hausdorffness, local compactness and first countability of manifolds.

In [13, Theorem 15] there is a proof that in a Tychonoff space 30 and 89 are equivalent.

The equivalence of condition 1 and conditions 98 to 101 is established in [12, Theorem 3.3].

The implication $1 \Rightarrow 103$ follows from 78 and [39, Theorem 8.17]. $103 \Rightarrow 104$ is trivial. $104 \Rightarrow 102$ is [34, Lemma 4.3]. $102 \Rightarrow 2$ is [34, Theorem 4.1].

Complete metrisability implies pseudocompleteness in any space and in turn pseudocompleteness implies α -favourability in a regular space, so $79 \Rightarrow 105 \Rightarrow 104$.

The implications $37 \Rightarrow 106$ and $106 \Rightarrow 28$ are shown in [10, Theorem 2.2].

The equivalence of 107 was already considered above in the context of 71.

Clearly every Baire space is Volterra and the converse holds in any locally convex topological vector space, [11, Theorem 3.4] so $107 \Leftrightarrow 108$. ■

4 Other properties of manifolds.

In this section we collect a few more properties which we may hope a manifold to possess.

Theorem 3 *Every manifold has a base of countable order.*

Proof: By [60, Theorem 2] every metric space has a base of countable order. As every manifold is locally metrisable it follows from [60, Theorem 1] that every manifold has a base of countable order. ■

Some standard conditions which manifolds may possess but which are weaker than metrisability are contained in the following theorem.

Theorem 4 *Suppose that the manifold M is metrisable. Then M is also normal, hereditarily normal, perfectly normal, separable, strongly hereditarily separable and has property (a).*

Proof outline: Every metrisable space is perfectly normal, normal and hereditarily normal. Every second countable space is separable and strongly hereditarily separable. Theorem 2(72) shows that metrisable manifolds satisfy property pp while in [26, Proposition 2.1] it is shown that every space having property pp has property (a). ■

There are manifolds which are normal but not metrisable, for example the long ray. The long ray also has property (a) (and, as shown in [26], even the stronger properties a-favourable and strongly a-favourable found in [42]).

The observant reader may have noticed that separability is absent as a criterion for metrisability in Theorem 2. The following example shows that it must be.

Example 5 *There is a manifold which is separable but not metrisable.*

One can make such a manifold out of the plane by replacing each point of the y -axis by an interval as follows. Let $S = \{(x, y) \in \mathbb{R}^2 / x \neq 0\}$ with the usual topology from \mathbb{R}^2 . Let $M = S \cup \{0\} \times \mathbb{R}^2$. For each $(0, \eta, \zeta) \in \mathbb{R}^3$ and each $r > 0$ let

$$W_{\eta, \zeta, r} = \{(x, y) \in S / \zeta - r < \frac{y - \eta}{|x|} < \zeta + r \text{ and } |x| < r\} \cup [\{0\} \times \{\eta\} \times (\zeta - r, \zeta + r)].$$

Topologise M by declaring $U \subset M$ to be open if and only if $U \cap S$ is open in S and for each $(0, \eta, \zeta) \in U \cap (M - S)$ there is $r > 0$ so that $W_{\eta, \zeta, r} \subset U$. Then M is a separable 2-manifold.

There are even manifolds which are both normal and separable but not metrisable, [56].

We need now some facts from Set Theory. The Continuum Hypothesis (CH), dating back to Cantor, states that any subset of \mathbb{R} either has the same cardinality as \mathbb{R} or is countable. Martin's Axiom (MA) can be expressed in various forms, the most topological of which is the following: in every compact, ccc, Hausdorff space the intersection of fewer than 2^{\aleph_0} dense open sets is dense. Recall the Baire Category Theorem which states that if X is Čech complete (ie X is a G_δ -set in βX ; for example every locally compact, Hausdorff space or every complete metric space) and $\{U_n / n \in \omega\}$ is a collection of open dense subsets of X then $\cap_{n \in \omega} U_n$ is dense in X . From the Baire Category theorem it is immediate that $\text{CH} \Rightarrow \text{MA}$. Both CH and MA are independent of the axioms of ZFC and otherwise of each other: thus there are models of Set Theory satisfying ZFC in which CH (and hence MA) holds, models in which MA holds but CH fails (denoted $\text{MA} + \neg \text{CH}$), and models in which MA (and hence CH) fails.

The question whether perfect normality is equivalent to metrisability for a manifold is an old one, dating back to [2]. It was shown in [55] that under $\text{MA} + \neg \text{CH}$ the two conditions are equivalent. On the other hand in [57] there is constructed an example of a perfectly normal non-metrisable manifold under CH. The same situation prevails when we consider strong hereditary separability. In [41] it is shown that under $\text{MA} + \neg \text{CH}$ every strongly hereditarily separable space is Lindelöf. On the other hand even when we combine the two notions the resulting manifold need not be metrisable in general; in [25] there is constructed under CH a manifold which is strongly hereditarily separable and perfectly normal but not metrisable. There are many other examples of conditions which are equivalent to metrisability for manifolds in some models of Set Theory but not equivalent in other models.

References

- [1] David F Addis and John H Gresham *A class of infinite-dimensional spaces. Part I: Dimension theory and Alexandroff's problem*, Fundamenta Mathematicae, 101(1978), 195–205.
- [2] P S Alexandroff, *On local properties of closed sets*, Annals of Mathematics, (2)26(1935), 1–35.
- [3] Richard F Arens, *A topology for spaces of transformations*, Annals of Mathematics, (2)47,1946), 480–495.
- [4] A V Arkhangel'skiĭ, *Hurewicz spaces, analytic sets, and fan tightness of function spaces*, Soviet Mathematics Doklady, 33(1986), 396–399.
- [5] A V Arhangel'skiĭ and R Z Buzyakova, *On linearly Lindelöf and strongly discretely Lindelöf spaces*, Topology Proceedings, 23(Summer 1998), 1–11.
- [6] Liljana Babinkostova, *Selective screenability game and covering dimension*, Topology Proceedings, 29(1)(2005), 13–17.
- [7] Z Balogh and M E Rudin, *Monotone Normality*, Topology and its Applications, 47(1992), 115–127.
- [8] Morton Brown, *The monotone union of open n -cells is an open n -cell*, Proceedings of the American Mathematical Society, 12(1961), 812–814.
- [9] G Cantor, *Ueber unendliche lineare Punktmannichfaltigkeiten*, Mathematische Annalen, 21(1883), 545–591.
- [10] Jiling Cao, David Gauld, Sina Greenwood and Abdul Mohamad, *Games and Metrisability of Manifolds*, New Zealand Journal of Mathematics, 37(2008), 1–8.
- [11] Jiling Cao and Heikki Junnila, *When is a Volterra space Baire?*, Topology and its Applications, 154(2007), 527–532.
- [12] Jiling Cao and Abdul Mohamad, *Metrizability, Manifolds and Hyperspace Topologies*, in preparation.
- [13] A Caserta, G Di Maio, Ljubiša D R Kočinac and Enrico Meccariello, *Applications of k -covers II*, Topology and its Applications, 153(2006), 3277–3293.
- [14] J Cheeger and J M Kister, *Counting Topological Manifolds*, Topology, 9(1970), 149–151.
- [15] C Costantini, L Holá and P Vitolo, *Tightness, character and related properties of hyperspace topologies*, Topology and its Applications, 142(2004), 245–292.
- [16] Satya Deo and David Gauld, *Boundedly Metacompact or Finitistic Spaces*, to appear.
- [17] G Di Maio, Ljubiša D R Kočinac and Enrico Meccariello, *Selection principles and hyperspace topologies*, Topology and its Applications, 153(2005), 912–923.
- [18] D L Fearnley, *Metrisation of Moore Spaces and Abstract Topological Manifolds*, Bulletin of the Australian Mathematical Society, 56(1997), 395–401.
- [19] J Fell, *A Hausdorff topology for the closed subsets of a locally compact non-Hausdorff space*, Proceedings of the American Mathematical Society, 13(1962), 472–476.

- [20] Otto Forster, *Lectures on Riemann Surfaces*, GTM 81 Springer-Verlag (1981).
- [21] M H Freedman and F Quinn, *Topology of 4-Manifolds*, Princeton University Press (1990).
- [22] P M Gartside and A M Mohamad, *Cleavability of Manifolds*, Topology Proceedings, 23(1998), 155–166.
- [23] P M Gartside and A M Mohamad, *Metrizability of Manifolds by Diagonal Properties*, Topology Proceedings, 24(1999), 621–640.
- [24] David Gauld, *Differential Topology: an introduction*, Lecture Notes in Mathematics 72(1982), Marcel Dekker. Reprint (2006), Dover.
- [25] David Gauld, *A strongly hereditarily separable, nonmetrizable manifold*, Topology and its Applications, 51(1993), 221–228.
- [26] David Gauld, *Covering Properties and Metrisation of Manifolds*, Topology Proceedings, 23(Summer 1998), 127–140.
- [27] David Gauld, *Some properties close to Lindelöf*, to appear.
- [28] David Gauld and Sina Greenwood, *Microbundles, Manifolds and Metrisability*, Proceedings of the American Mathematical Society, 128 (2000), 2801–2807.
- [29] David Gauld, Sina Greenwood and Zbigniew Piotrowski, *On Volterra spaces III: Topological operations*, Topology Proceedings, 23(1998), 167–182.
- [30] David Gauld and Frédéric Mynard, *Metrisability of Manifolds in Terms of Function Spaces*, Houston Journal of Mathematics, 31(2005), 199–214.
- [31] David Gauld and M K Vamanamurthy, *Covering Properties and Metrisation of Manifolds 2*, Topology Proceedings, 24(Summer 1999), 173–185.
- [32] E. Grabner, G. Grabner and J. E. Vaughan, *Nearly metacompact spaces*, Topology and its Applications, 98(1999), 191–201.
- [33] Gary Gruenhage, *Generalized Metric Spaces*, in K Kunen and J Vaughan, eds, “Handbook of Set-Theoretic Topology” (Elsevier, 1984), 423–501.
- [34] Gary Gruenhage, *The Story of a Topological Game*, Rocky Mountain Journal of Mathematics, 36(2006), 1885–1914.
- [35] Gary Gruenhage and Daniel K Ma, *Baireness of $C_k(X)$ for locally compact X* , Topology and its Applications, **80** (1997), 131–139.
- [36] R W Heath, D J Lutzer and P L Zenor, *Monotonically Normal Spaces*, Transactions of the American Mathematical Society, 178(1973), 481–493.
- [37] John G Hocking and Gail S Young, *Topology*, Addison-Wesley (1961).
- [38] I M James, *Topological and Uniform Structures*, Undergraduate Texts in Mathematics (1987), Springer Verlag.
- [39] A. S. Kechris, *Classical Descriptive Set Theory*, Springer-Verlag, New York, 1995.
- [40] J M Kister, *Microbundles are Fibre Bundles*, Annals of Mathematics, (2) 80 (1964), 190–199.

- [41] K Kunen, *Strong S and L spaces under MA* , in G M Reed, ed, “Set-Theoretic Topology” (Academic Press, New York, 1977), 265–268.
- [42] M V Matveev, *Some Questions on Property (a)*, Questions and Answers in General Topology, 15(1997), 103–111.
- [43] Robert A McCoy and Ibula Ntantu, *Topological Properties of Spaces of Continuous Functions*, Lecture Notes in Mathematics 1315, Springer-Verlag, 1988.
- [44] J Milnor, *Lectures on the h -Cobordism Theorem*, Princeton Mathematical Notes (1965), Princeton University Press.
- [45] A M Mohamad, *Metrization and semimetrization theorems with applications to manifolds*, Acta Mathematica Hungarica, 83 (4) (1999), 383–394.
- [46] Frédéric Mynard, *First countability, sequentiality and tightness of the upper Kuratowski convergence*, preprint series. LAAO, Université de Bourgogne, Octobre 2000.
- [47] Peter Nyikos, *The Theory of Nonmetrizable Manifolds*, in K Kunen and J Vaughan, eds, “Handbook of Set-Theoretic Topology” (Elsevier, 1984), 634–684.
- [48] Peter Nyikos, *Various Smoothings of the Long Line and Their Tangent Bundles*, Advances in Mathematics, 93(1992), 129–213.
- [49] Peter Nyikos, *Mary Ellen Rudin’s Contributions to the Theory of Nonmetrizable Manifolds*, in F D Tall, ed, “The Work of Mary Ellen Rudin” (Annals of the New York Academy of Sciences, 705(1993)), 92–113.
- [50] A R Pears, *Dimension Theory of General Spaces*, Cambridge University Press, 1975.
- [51] H Poincaré, *Second complement a l’analyse situs*, Proceedings of the London Mathematical Society, 32(1900), 277–302.
- [52] H Poincaré, *Cinquieme complement a l’analyse situs*, Rendiconti Circolo Matematici Palermo, 18(1904), 45–110.
- [53] G M Reed and P L Zenor, *A metrization theorem for normal Moore spaces*, in N M Stavrakas and K R Allen, eds, “Studies in Topology” (Academic Press, New York, 1974), 485–488.
- [54] G M Reed and P L Zenor, *Metrization of Moore spaces and generalized manifolds*, Fundamenta Mathematicae, 91(1976), 203–210.
- [55] M E Rudin, *The undecidability of the existence of a perfectly normal non-metrizable manifold*, Houston Journal of Mathematics, 5(1979), 249–252.
- [56] M E Rudin, *Two Nonmetrizable Manifolds*, Topology and its Applications, 35(1990), 137–152.
- [57] M E Rudin and P L Zenor, *A perfectly normal non-metrizable manifold*, Houston Journal of Mathematics, 2(1976), 129–134.
- [58] Yu M Smirnov, *On Metrization of Topological Spaces*, Uspehi Matematiki Nauk, 6(1951), #6(46), 100–111 (American Mathematical Society Translations #91).

- [59] M G Tkačenko, *Ob odnom svoistve bicomactov (On a property of compact spaces)*, Seminar po obshchei topologii (A seminar on general topology), Moscow State University P. H., Moscow (1981), 149–156 (Russian).
- [60] J M Worrell Jr and H H Wicke, *Characterizations of Developable Topological Spaces*, Canadian Journal of Mathematics, 17(1965), 820–830.
- [61] Scott W Williams and Haoxuan Zhou, *Strong Versions of Normality*, General Topology and its Applications, Proceedings of the 5th NE Conference, New York 1989, in Lecture Notes in Pure and Applied Mathematics, 134 (Marcel Dekker, New York, 1991), 379–389.

d.gauld@auckland.ac.nz

The Department of Mathematics
The University of Auckland
Private Bag 92019
Auckland
New Zealand.